MEDICINE INFORMATION

Docetaxel and carboplatin

A guide to docetaxel and carboplatin, with practical information about the steroid, blood tests and symptoms to report.

At a glance

Treatment	Docetaxel + Carboplatin
When it may be used	The combination is one chemotherapy option for some people with ovarian, fallopian tube or primary peritoneal cancer.
How it is given	Docetaxel and carboplatin are both given by drip into a vein.
Before you start	Tell your team about kidney or liver problems, diabetes, past drip reactions and any tingling or numbness.

What this treatment is and how it works

Docetaxel stops cells dividing normally. Carboplatin damages the genetic material in cells. The drugs work in different ways and both can lower blood counts.

When it may be used

Treatment calendars and reasons for choosing a taxane vary. Ask why docetaxel is recommended for you.

Before treatment

If you are given steroid tablets before treatment, ask which doses to take at home.

How treatment is given

Both medicines are given through a drip in hospital. You may receive medicines for sickness and to reduce reactions to docetaxel. The team will explain whether a white-cell support injection is part of your plan.

Tests and monitoring during treatment

Blood counts and kidney function help your team plan treatment. Liver tests also matter for docetaxel. Tell the team about changes in walking, writing, grip or using buttons, so they can assess any nerve symptoms.

Side effects

Common or more frequent side effects

- **Changes in blood counts.** Treatment can lower the white cells that fight infection, the red cells that carry oxygen and the platelets that stop bleeding. You may feel tired or get breathless sooner than usual. Let the unit know about nosebleeds, new bruises or small red or purple spots. Fever or shaking chills needs a call to the urgent number even if your next blood test is already booked.
- **Nail changes and hair loss.** Nails may discolour, become tender or loosen. Keep them clean and tell the team about pain, swelling or discharge, especially if your blood counts are low. Hair loss can start during treatment. Ask early about scalp cooling where appropriate, because it needs to be arranged before chemotherapy.
- **Fluid retention.** Docetaxel can cause swelling of the ankles, legs or fingers, sometimes after several cycles. Tell the team about swelling that increases, rapid weight gain or becoming more breathless with your usual activities. Follow the prescribed steroid schedule, which may help reduce fluid retention as well as infusion reactions.
- **Tingling, altered feeling or pain in hands and feet.** Give examples of what has changed, such as difficulty with buttons, holding a cup or getting your balance on steps. Tell the team if the sensation lasts until the next visit. If temperature is hard to judge with your fingers or toes, take care around hot water and heating pads.

Other reported side effects

Tell your team about ringing in the ears or hearing changes.

Serious effects to recognise

- **Infusion reaction.** A reaction can develop even after you have received platinum chemotherapy before. Tell the nurse immediately about sudden itching, rash, dizziness, back pain, chest tightness or breathing changes. Keep information about past reactions so staff can plan future infusions safely.

When to seek urgent help

Call your cancer team straight away if you have fever, shaking chills, unexpected bleeding, new breathlessness or cannot keep fluids down. Follow your hospital’s written temperature instructions. If you have no written threshold, call at 37.5°C (99.5°F) or above, or below 36°C (96.8°F). Call if you feel very unwell even when your temperature is normal.

Seek emergency care for severe breathing difficulty, swelling of the face or throat, chest pain, collapse or bleeding that will not stop.

Practical advice for this treatment

Before the first day

Check the times of any steroid tablets prescribed before docetaxel. Tell the nurse if you could not take them. If you have diabetes, ask whether you need extra glucose checks while taking the steroid and whom to call about a high result. Bring the names of previous platinum treatments and any reactions.

Changes around the nails, feet and ankles

Docetaxel can cause tender nails, changes in nail colour and swelling. Keep nails clean and protect them from knocks. Tell the team about redness, discharge, swelling or nail pain that makes walking or using your hands difficult. Ankle swelling can develop over several cycles. Tell the team if it increases or if you also feel breathless.

Between visits

A low blood count can develop after you feel you have recovered from the infusion. Keep a thermometer where you can find it. Contact the unit for fever or feeling suddenly unwell. Keep track of unusual bruises, nosebleeds and how often they occur.

Medicines, interactions and vaccinations

Ask your pharmacist to check prescriptions, supplements and other remedies. If you have diabetes, ask how steroid days may affect your blood sugar. Check before live vaccines.

Everyday life

Bring the steroid instructions and allow for extra time at your first visit. Ask about managing taste changes, getting help with transport and balancing activity with rest.

Pregnancy, fertility and breastfeeding

Discuss contraception, fertility and breastfeeding before treatment. Ask when precautions end for each medicine.

After treatment

Ask how the team will assess response and what the next step will be. Report tingling, swelling, hearing changes or unusual tiredness if they continue after treatment.

Frequently asked questions

What are the steroid tablets for?

They help lower the chance of some docetaxel reactions and fluid retention.

What if I forget the steroid?

Call the treatment unit before coming in and tell them which doses you missed.

Why are kidney tests needed?

The kidneys clear carboplatin. Results help the team decide how much can be given safely.

Is this the same as carboplatin and paclitaxel?

The platinum drug is the same. Docetaxel and paclitaxel are different chemotherapy medicines.

Can I react after several uneventful visits?

Yes. Tell the unit if you have received carboplatin before and whether you had any reaction. Report sudden itching, flushing, tightness in the chest or throat, dizziness or breathlessness immediately during a drip.

Can nail changes be helped?

Tell the team about pain, lifting or signs of infection. Ask what nail care is suitable.

Should I report swelling?

Yes. New swelling, quick weight gain or new breathlessness should be checked.

Can I ask about scalp cooling?

Ask before the first cycle whether it is suitable and available.

Individual medicine guides

- Docetaxel
- Carboplatin

Sources and further reading

Read this guide alongside your hospital's plan. Ask your pharmacist to confirm the exact product dispensed in India and follow the instructions for that product.

Summary of Product Characteristics (SmPC) and prescribing information

- Docetaxel (Taxotere) - EU product page with SmPC
- Carboplatin infusion - UK SmPC

Clinical protocols and other references

- eviQ docetaxel and carboplatin protocol in ovarian cancer

Additional sources

- eviQ patient information for carboplatin and docetaxel

Editorially approved by Dr (Prof) Vijay Agarwal

Professor and Lead Consultant - Medical Oncology

Review record updated: 14 November 2026 · Editorial review policy